

Carolyn D. Ashworth, M.D.

Chandra Garvey, R.N.C.P.N.P.

Michele Hickman, R.N.C.P.N.P.

Wt. _____ % Ht. _____ % Head _____ %

TWO-MONTH VISIT

Prevent falls; never leave baby alone on beds, couches, etc.

Do not drink hot liquids while holding the baby; no smoking in baby's environment.

Do not start solids or add cereal to formula unless advised to by health care provider.

Breast-feeding: Feed 8-12 times a day, growth spurt may occur around 2-1/2 months, and baby may need to be fed more frequently.

Bottle feeding: Never prop bottle.

Put baby in crib when drowsy, but before he/she is completely asleep.

Try to establish a regular naptime in the morning and afternoon.

Best stimulation for baby is playing with, talking to and cuddling.

Watch for signs of illness: Fever, poor feeding, decreased activity, vomiting, diarrhea; Call office if temperature is > 101degrees, if fewer than 6 wet diapers/day, or if you are concerned about your baby's symptoms/behavior.

AT THIS VISIT YOUR CHILD MAY RECEIVE THE FOLLOWING IMMUNIZATIONS:

DTaP Hib Polio Prevnar Hep B

NEXT WELL CHILD VISIT:

At 4 months of age: 2nd DTaP, Hib, Polio, and Prevnar will be given.